



Tehran University of Medical Sciences
School of Nursing and Midwifery

**A survey of Causes of Failure to Receive Prenatal Care and
Associated Factors in Delivered Women in Iranian Hospital- Dubai
2013**

A Thesis submitted to the Graduate Studies Office in partial fulfillment of the requirements for The degree
of Master in Midwifery
(branch of Maternal and Child Health)

Student: Mohammad Ibrahim Fatima

Supervisor:
Golian Tehrani Shahnaz, MSc

Advisor:
Pakgozar Menoo, PhD

Statistics Advisor:
Haghani Hamed, PhD

2013-2014

Abstract

Background: Reproductive health is one of the essential prerequisites for achieving sustainable development and in contemporary societies is considered. The objective of the present study was to evaluate the effective factors in prenatal care.

Methods: A cross-section study was performed on 520 pregnant women who had delivered in Iranian Red Crescent Hospital-Dubai. Data collected using a questionnaire that included demographic characteristics, referring to the lack of prenatal care and refer pregnant women to get the serviced one. After authenticating the scientific literature and the opinion of the questionnaire teachers using SPSS software and statistical methods of analysis received. Based on the results obtained in terms of age at first pregnancy ($p=0.01$), Gestational age at birth ($p=0.01$), mother's education, husband's education ($p=0.00$), mother's occupation, husband's occupation ($p=0.00$), socioeconomic status ($p=0.002$), insurance status ($p=0.00$), religion ($p=0.017$), pregnancy ($p=0.01$) had assign if I cant But in terms of nationality ($P =0.606$), infertility ($P =0.643$), abortion ($P =0.597$), preterm delivery ($P =0.211$), living child ($P =0.46$), son of the dead ($P =0.597$) difference not significant. In general it can be concluded that the age at first pregnancy, gestational age at delivery, maternal education level and husbands mother and wife, socioeconomic status, insurance status, religion, unwanted pregnancy are factors that play a fundamental role in the lack of care for mothers during pregnancy and the risk of these people are Roper.

Key Words: Prenatal care ,pregnant women, Effective Factors